

## Ph 203. Solutions HW #1

1.) Energy levels.

a) Infinite potential well

$$V = \begin{cases} \infty & \text{for } r > a \\ 0 & \text{for } r < a \end{cases}$$

Spherically symmetric potential, so can write the total wavefunction as

$$\Psi_{klm}(r, \theta, \phi) = R_{kl}(r)Y_m^l(\theta, \phi),$$

with the radial part  $R_{kl}(r) = \frac{u(r)}{r}$ .  
Schrödinger equation for  $r < a$ :

$$-\frac{1}{2\mu} \frac{\partial^2 u(r)}{\partial r^2} + \frac{l(l+1)}{2\mu r^2} u(r) = \frac{k^2}{2\mu} u(r),$$

with energy  $E = \frac{k^2}{2\mu}$ .

The solutions are spherical Bessel functions  $j_l(kr)$  and energy levels are determined by  $j_l(ka) = 0$ .


Roots of  $j_l(x) = 0$  are:

	n=1	n=2	n=3
$l = 0$	3.14	6.28	9.42
$l = 1$	4.49	7.72	
$l = 2$	5.76	9.10	
$l = 3$	7.00		
$l = 4$	8.18		
$l = 5$	9.36		

#fermions( $s = 1/2$ ) =  $2 \times$  degeneracy

Energy levels, labeled by  $nl$ (degeneracy), (from lowest to highest):

1s(1), 1p(3), 1d(5), 2s(1), 1f(7), 2p(3), 1g(9), 2d(5), 1h(11), 3s(1)



b) Radial SHO: use separation of variables in cartesian coordinates

$$\begin{aligned}
 H &= \omega \left( n_x + \frac{1}{2} \right) + \omega \left( n_y + \frac{1}{2} \right) + \omega \left( n_z + \frac{1}{2} \right) \\
 &= \omega \left( n_x + n_y + n_z + \frac{3}{2} \right) = \omega \left( n + \frac{3}{2} \right),
 \end{aligned}$$


where  $n = n_x + n_y + n_z$ .

Energy levels (from lowest) labeled by  $N = n + 1$ :

N(degeneracy)	$l$
1(1)	0
2(3)	1
3(6)	0, 2
4(10)	1, 3
5(15)	0, 2, 4

Degeneracy - count all equivalent ways one can choose  $n_x, n_y, n_z$  to get  $n$

Angular momentum:  $l = N - 1, N - 3, \dots$



2.) Spin projection

$$\hat{S}_{\hat{n}} = \frac{1}{2} \vec{\sigma} \cdot \hat{n} = \frac{1}{2} \begin{pmatrix} \cos \theta & \sin \theta e^{-i\phi} \\ \sin \theta e^{i\phi} & -\cos \theta \end{pmatrix},$$

where  $\hat{n} = (\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta)$ . Eigenvalues:  $\lambda = \pm \frac{1}{2}$ . For  $\lambda = \frac{1}{2}$ , eigenvector is  $\chi = (\cos \frac{\theta}{2}, \sin \frac{\theta}{2} e^{i\phi})$ .

$$\Rightarrow a = \cos \frac{\theta}{2}, b = \sin \frac{\theta}{2} e^{i\phi}$$

\*Answers may differ depending on phase rotations\*

3.) Adding three spin 1/2 particles

Symmetric under exchange

$$\begin{aligned} \left| \frac{3}{2}, \frac{3}{2} \right\rangle &= \uparrow\uparrow\uparrow \\ \left| \frac{3}{2}, \frac{1}{2} \right\rangle &= \frac{1}{\sqrt{3}}(\uparrow\uparrow\downarrow + \uparrow\downarrow\uparrow + \downarrow\uparrow\uparrow) \\ \left| \frac{3}{2}, -\frac{1}{2} \right\rangle &= \frac{1}{\sqrt{3}}(\uparrow\downarrow\downarrow + \downarrow\uparrow\downarrow + \downarrow\downarrow\uparrow) \\ \left| \frac{3}{2}, -\frac{3}{2} \right\rangle &= \downarrow\downarrow\downarrow \end{aligned}$$

Symmetric under  $1 \leftrightarrow 2$

$$\begin{aligned} \left| \frac{1}{2}, \frac{1}{2} \right\rangle &= \sqrt{\frac{2}{3}} \uparrow\uparrow\downarrow - \frac{1}{\sqrt{6}}(\uparrow\downarrow\uparrow + \downarrow\uparrow\uparrow) \\ \left| \frac{1}{2}, -\frac{1}{2} \right\rangle &= -\sqrt{\frac{2}{3}} \downarrow\downarrow\uparrow + \frac{1}{\sqrt{6}}(\uparrow\downarrow\downarrow + \downarrow\uparrow\downarrow) \end{aligned}$$

Antisymmetric under  $1 \leftrightarrow 2$

$$\begin{aligned} \left| \frac{1}{2}, \frac{1}{2} \right\rangle &= \frac{1}{\sqrt{2}}(\uparrow\downarrow\uparrow - \downarrow\uparrow\uparrow) \\ \left| \frac{1}{2}, -\frac{1}{2} \right\rangle &= \frac{1}{\sqrt{2}}(\uparrow\downarrow\downarrow - \downarrow\uparrow\downarrow) \end{aligned}$$

4.) Bertulani 1.6

Incoming proton  $p_1 = (E, 0, 0, p)$

proton at rest  $p_2 = (m_p, 0, 0, 0)$

$$\Rightarrow s = (p_1 + p_2)^2 = 2m_p(m_p + E)$$


Final state particles: min  $E$  when final particles at rest in C.M. frame

$$\sum_{final} p'_i = (4m_p, 0, 0, 0)$$

$$\Rightarrow s = \left( \sum_{final} p'_i \right)^2 = 16m_p^2$$

Equating  $s$ , we get  $E = 7m_p$ , so kinetic energy is  $KE = 6m_p \approx 5.6$  GeV

5.) Parity of the pion



**Initial state**

Pion is spin 0 and  $\pi^-$  captured at rest  $\Rightarrow l = 0$ .

Parity of the initial state =  $P_\pi(-1)^l P_d = P_\pi$  for  $l = 0, P_d = +1$ .

Deuteron is spin 1.

Total angular momentum of the initial state:  $J = 1$

**Final state**

n-n has overall antisymmetric wavefunction and total angular momentum  $J = 1$ .

Possibilities ( $s$  is total spin,  $l$  is orbital angular momentum):

[1]  $s = 0, l = 0 \rightarrow J = 0$ , not allowed

[2]  $s = 0, l = 1 \rightarrow$  symmetric wavefunction, not allowed

[3]  $s = 1, l = 0 \rightarrow$  symmetric wavefunction, not allowed

[4]  $s = 1, l = 1$ , can have:

○  $J = 0 \rightarrow$  symmetric wavefunction, not allowed

○  $J = 1 \rightarrow$  antisymmetric wavefunction

○  $J = 2 \rightarrow$  symmetric wavefunction, not allowed

Only the  $J = 1, s = 1, l = 1$  option gives a totally antisymmetric state with  $J = 1$ . Since  $l = 1$ , the parity of the final state is  $P_{2n} = -1$ . Since parity is conserved in this process, we deduce  $P_\pi = -1$ .