

# Homework Set #1

\*Problem 1.5 Consider the wave function

$$\Psi(x, t) = Ae^{-\lambda|x|}e^{-i\omega t},$$

where  $A$ ,  $\lambda$ , and  $\omega$  are positive real constants. (We'll see in Chapter 2 what potential ( $V$ ) actually produces such a wave function.)

- (a) Normalize  $\Psi$ .
- (b) Determine the expectation values of  $x$  and  $x^2$ .
- (c) Find the standard deviation of  $x$ . Sketch the graph of  $|\Psi|^2$ , as a function of  $x$ , and mark the points  $(\langle x \rangle + \sigma)$  and  $(\langle x \rangle - \sigma)$ , to illustrate the sense in which  $\sigma$  represents the "spread" in  $x$ . What is the probability that the particle would be found outside this range?

See Additional Problem on Next Page

**Problem 1.17** A particle is represented (at time  $t = 0$ ) by the wave function

$$\Psi(x, 0) = \begin{cases} A(a^2 - x^2), & \text{if } -a \leq x \leq +a, \\ 0, & \text{otherwise.} \end{cases}$$

- (a) Determine the normalization constant  $A$ .
- (b) What is the expectation value of  $x$  (at time  $t = 0$ )?
- (c) What is the expectation value of  $p$  (at time  $t = 0$ )? (Note that you *cannot* get it from  $p = md\langle x \rangle/dt$ . Why not?)
- (d) Find the expectation value of  $x^2$ .
- (e) Find the expectation value of  $p^2$ .
- (f) Find the uncertainty in  $x$  ( $\sigma_x$ ).
- (g) Find the uncertainty in  $p$  ( $\sigma_p$ ).
- (h) Check that your results are consistent with the uncertainty principle.